

$$\begin{aligned}
 1. \quad & \frac{2\sqrt{2} \cdot 3\sqrt{4}}{3\sqrt{2} \cdot \sqrt{8}} = \frac{2 \cdot 3\sqrt{2^3 \cdot 4}}{3\sqrt{2} \cdot 2\sqrt{2^3}} \\
 & = \frac{6\sqrt{2^5}}{3\sqrt{2} \cdot 2\sqrt{2^3}} \\
 & = \frac{2^{\frac{5}{6}}}{\frac{1}{2^3} \cdot \frac{3}{2^2}} \\
 & = \frac{2^{\frac{5}{6}}}{\frac{11}{2^6}} \\
 & = 2^{\frac{5}{6} - \frac{11}{6}} \\
 & = 2^{-1} = \frac{1}{2}
 \end{aligned}$$

Cevap: E

$$\begin{aligned}
 2. \quad & (\sqrt{2} - \sqrt{5})^3 = (\sqrt{2})^3 - 3(\sqrt{2})^2 \cdot \sqrt{5} + 3 \cdot \sqrt{2} \cdot (\sqrt{5})^2 - (\sqrt{5})^3 \\
 & = 2\sqrt{2} - 3 \cdot 2 \cdot \sqrt{5} + 3 \cdot 5 \cdot \sqrt{2} - 5\sqrt{5} \\
 & = 2\sqrt{2} - 6\sqrt{5} + 15\sqrt{2} - 5\sqrt{5} \\
 & = 17\sqrt{2} - 11\sqrt{5}
 \end{aligned}$$

Cevap: B

$$3. \quad x \in \mathbb{R}^+$$

$$\begin{aligned}
 & \frac{4\sqrt{x^3} \cdot 3\sqrt{x^4}}{12\sqrt{x^{24}}} = \frac{x^{\frac{3}{4}} \cdot x^{\frac{4}{3}}}{\frac{24}{x^{12}}} \\
 & = \frac{x^{\frac{3}{4} + \frac{4}{3}}}{\frac{24}{x^{12}}} \\
 & = x^{\frac{25}{12} - \frac{24}{12}} \\
 & = x^{\frac{1}{12}} = \sqrt[12]{x}
 \end{aligned}$$

Cevap: D

$$\begin{aligned}
 4. \quad & \frac{(3\sqrt{2})^2 - (2\sqrt{3})^2}{\frac{3}{2^2} - \frac{1}{2^2}} = \frac{9 \cdot 2 - 4 \cdot 3}{\sqrt{2^3} - \sqrt{2}} \\
 & = \frac{18 - 12}{\sqrt{8} - \sqrt{2}} \\
 & = \frac{6}{2\sqrt{2} - \sqrt{2}} \\
 & = \frac{6}{\frac{\sqrt{2}}{(\sqrt{2})}} \\
 & = \frac{6\sqrt{2}}{2} = 3\sqrt{2}
 \end{aligned}$$

Cevap: E

$$5. \quad \sqrt{4+\sqrt{7}} - \sqrt{4-\sqrt{7}} = x \text{ olsun.}$$

Eşitliğin her iki tarafının karesini alalım.

$$\begin{aligned}
 & (\sqrt{4+\sqrt{7}} - \sqrt{4-\sqrt{7}})^2 = x^2 \\
 & 4 + \sqrt{7} - 2 \cdot \underbrace{\sqrt{(4+\sqrt{7})(4-\sqrt{7})}}_{\text{İki kare farkı}} + 4 - \sqrt{7} = x^2 \\
 & 8 - 2 \cdot \sqrt{4^2 - (\sqrt{7})^2} = 8 - 2 \cdot \sqrt{16 - 7} = 8 - 2 \cdot \sqrt{9} \\
 & = 8 - 2 \cdot 3 \\
 & = 8 - 6 = 2 = x^2 \\
 & \Rightarrow x = \sqrt{2}
 \end{aligned}$$

Cevap: D

$$\begin{aligned}
 6. \quad & \sqrt{13 - 2\sqrt{30}} + \sqrt{5 - 2\sqrt{6}} + \sqrt{2} \\
 & \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 & 10+3 \quad 10 \cdot 3 \quad 3+2 \quad 3 \cdot 2 \\
 & = \sqrt{10} - \sqrt{3} + \sqrt{3} - \sqrt{2} + \sqrt{2} = \sqrt{10}
 \end{aligned}$$

Cevap: A

$$7. \frac{\sqrt{3,61} + \sqrt{1,69}}{\sqrt{1,44} - \sqrt{0,64}} = \frac{\sqrt{\frac{361}{100}} + \sqrt{\frac{169}{100}}}{\sqrt{\frac{144}{100}} - \sqrt{\frac{64}{100}}} \\ = \frac{\frac{19}{10} + \frac{13}{10}}{\frac{12}{10} - \frac{8}{10}} \\ = \frac{\frac{32}{10}}{\frac{4}{10}} = 8$$

Cevap: A

$$8. \sqrt{0,006} \cdot \sqrt{0,08} \cdot \sqrt{0,3} = \sqrt{\frac{6}{1000} \cdot \frac{8}{100} \cdot \frac{3}{10}} \\ = \sqrt{\frac{144}{1000000}} \\ = \frac{12}{1000} \\ = 0,012$$

$$9. \sqrt{3 + \sqrt{2\sqrt{7} + 1}} - \sqrt{3 - \sqrt{2\sqrt{7} + 1}} = x \text{ olsun.}$$

Eşitliğin her iki tarafının karesini alalım.

$$(\sqrt{3 + \sqrt{2\sqrt{7} + 1}} - \sqrt{3 - \sqrt{2\sqrt{7} + 1}})^2 = x^2$$

$$3 + \sqrt{2\sqrt{7} + 1} - 2 \cdot \sqrt{9 - (2\sqrt{7} + 1)} + 3 - \sqrt{2\sqrt{7} + 1} = x^2$$

$$6 - 2 \cdot \sqrt{9 - 2\sqrt{7} - 1} = x^2$$

$$6 - 2 \cdot \sqrt{8 - 2\sqrt{7}} = x^2$$

↓ ↓
7+1 7.1

$$6 - 2 \cdot (\sqrt{7} - 1) = x^2$$

$$6 - 2\sqrt{7} + 2 = x^2$$

$$8 - 2\sqrt{7} = x^2$$

$$\sqrt{8 - 2\sqrt{7}} = x \Rightarrow x = \sqrt{7} - 1$$

↓ ↓
7+1 7.1

TASARI EĞİTİM YAYINLARI

$$10. \sqrt[3]{2 \cdot \sqrt{21 - \sqrt{23 + \sqrt{4}}}} = \sqrt[3]{2 \cdot \sqrt{21 - \sqrt{23 + 2}}} \\ = \sqrt[3]{2 \cdot \sqrt{21 - \sqrt{25}}} \\ = \sqrt[3]{2 \cdot \sqrt{21 - 5}} \\ = \sqrt[3]{2 \cdot \sqrt{16}} \\ = \sqrt[3]{2 \cdot 4} = \sqrt[3]{8} = \sqrt[3]{2^3} \\ = 2$$

Cevap: E

$$11. \sqrt{\frac{27}{4}} + \sqrt{\frac{3}{4}} - \sqrt{\frac{75}{4}} = \frac{\sqrt{9 \cdot 3}}{2} + \frac{\sqrt{3}}{2} - \frac{\sqrt{25 \cdot 3}}{2} \\ = \frac{3\sqrt{3} + \sqrt{3} - 5\sqrt{3}}{2} \\ = \frac{-\sqrt{3}}{2}$$

Cevap: D

$$12. \frac{3x - 2}{\sqrt{3} + \sqrt{2}} = \frac{\sqrt{5 - 2\sqrt{6}}}{\sqrt{3+2} \quad \sqrt{3 \cdot 2}} \Rightarrow \frac{3x - 2}{\sqrt{3} + \sqrt{2}} = \sqrt{3} - \sqrt{2}$$

Cevap: D

$$\Rightarrow 3x - 2 = (\underbrace{(\sqrt{3} - \sqrt{2}) \cdot (\sqrt{3} + \sqrt{2})}_{\text{iki kare farkı}})$$

$$\Rightarrow 3x - 2 = (\sqrt{3})^2 - (\sqrt{2})^2$$

$$\Rightarrow 3x - 2 = 3 - 2 = 1$$

$$\Rightarrow 3x = 3 \Rightarrow x = 1$$

Cevap: E

$$13. \frac{\sqrt{0,98} + \sqrt{1,62} + \sqrt{0,72}}{\sqrt{2,42}} \\ = \frac{\sqrt{\frac{98}{100}} + \sqrt{\frac{162}{100}} + \sqrt{\frac{72}{100}}}{\sqrt{\frac{242}{100}}} \\ = \frac{\frac{\sqrt{49 \cdot 2}}{10} + \frac{\sqrt{81 \cdot 2}}{10} + \frac{\sqrt{36 \cdot 2}}{10}}{\sqrt{121 \cdot 2}} \\ = \frac{7\sqrt{2} + 9\sqrt{2} + 6\sqrt{2}}{10} \cdot \frac{10}{11\sqrt{2}} = \frac{22\sqrt{2}}{11\sqrt{2}} = 2$$

Cevap: C

Cevap: C

14. $\frac{1+\sqrt{3}}{\sqrt{5}-\sqrt{3}} = a \Rightarrow \frac{1}{a} = \frac{\sqrt{5}-\sqrt{3}}{\sqrt{3}+1}$
 $\frac{\sqrt{3}+\sqrt{5}}{\sqrt{3}-1} = b \text{ olsun. } \Rightarrow x \frac{b}{\sqrt{3}-1} = \frac{\sqrt{5}+\sqrt{3}}{\sqrt{3}-1}$

$$\frac{1}{a} \cdot b = \frac{\sqrt{5}-\sqrt{3}}{\sqrt{3}+1} \cdot \frac{\sqrt{5}+\sqrt{3}}{\sqrt{3}-1}$$

$$\frac{b}{a} = \frac{(\sqrt{5})^2 - (\sqrt{3})^2}{(\sqrt{3})^2 - 1^2}$$

$$\frac{b}{a} = \frac{5-3}{3-1}$$

$$\frac{b}{a} = \frac{2}{2} = 1 \Rightarrow b = a$$

Cevap: B

15. $\sqrt{\frac{10}{9} - \sqrt{\frac{10}{9} - \sqrt{\frac{10}{9} \dots}}} = ?$

$a(a+1) = \frac{10}{9}$ olsun. Buradan $a = \frac{2}{3}$, $a+1 = \frac{5}{3}$ olur.

O halde verilen çıkarma işleminin sonucu $\boxed{a = \frac{2}{3}}$ 'tür.

(Küçük olan sayı)

Cevap: D

16. $\sqrt{\frac{abc,abc}{a,bcabc}} + \sqrt{\frac{1111}{11} + 20}$
 $= \sqrt{\frac{abcabc00}{abcabc}} + \sqrt{\frac{1111+220}{11}}$
 $= \sqrt{100} + \sqrt{\frac{1331}{11}} = 10 + \sqrt{121} = 10 + 11 = 21$

Cevap: E

17. $4\sqrt{(1-\sqrt{3})^4} + 3\sqrt[3]{(1-\sqrt{3})^3}$
 $= \underbrace{|1-\sqrt{3}|}_{<0} + (1-\sqrt{3})$
 $= -(1-\sqrt{3}) + (1-\sqrt{3})$
 $= -\cancel{1} + \cancel{\sqrt{3}} + \cancel{1} - \cancel{\sqrt{3}} = 0$

Cevap: A

18. $\frac{\frac{4}{\sqrt{3}-1} + \frac{2}{\sqrt{2}+1}}{\frac{1}{\sqrt{5}-2\sqrt{6}}} = \frac{\frac{2}{\cancel{2}} \cdot (\sqrt{3}+1) + \frac{2 \cdot (\sqrt{2}-1)}{\cancel{2}}}{\frac{1}{\sqrt{3}-\sqrt{2}}}$
 $\downarrow \quad \downarrow$
 $3+2 \quad 3 \cdot 2$
 $= (2\sqrt{3} + 2 + 2\sqrt{2} - 2) \cdot \frac{(\sqrt{3}-\sqrt{2})}{1}$
 $= 2(\sqrt{3} + \sqrt{2})(\sqrt{3} - \sqrt{2})$
 $= 2((\sqrt{3})^2 - (\sqrt{2})^2) = 2(3-2) = 2$

Cevap: E

19. $\frac{2}{\sqrt{3}-\sqrt{2}} + \frac{1}{3+\sqrt{8}} - \frac{2}{2-\sqrt{3}}$
 $= \frac{2(\sqrt{3}+\sqrt{2})}{(\sqrt{3})^2 - (\sqrt{2})^2} + \frac{3-\sqrt{8}}{3^2 - (\sqrt{8})^2} - \frac{2 \cdot (2+\sqrt{3})}{2^2 - (\sqrt{3})^2}$
 $= \frac{2\sqrt{3}+2\sqrt{2}}{3-2} + \frac{3-\sqrt{8}}{9-8} - \frac{4+2\sqrt{3}}{4-3}$
 $= 2\sqrt{3} + 2\sqrt{2} + 3 - 2\sqrt{2} - 4 - 2\sqrt{3} = 3 - 4 = -1$

Cevap: E

20. $a = \sqrt{(1-\sqrt{5})^2} = \underbrace{|1-\sqrt{5}|}_{<0} = -1 + \sqrt{5}$
 $b = \sqrt{(\sqrt{5}-\sqrt{20})^2} = \sqrt{(\sqrt{5}-\sqrt{4 \cdot 5})^2}$
 $= \sqrt{(\sqrt{5}-2\sqrt{5})^2} = \sqrt{(-\sqrt{5})^2} = \sqrt{5}$
 $\Rightarrow a+b = -1 + \sqrt{5} + \sqrt{5} = 2\sqrt{5} - 1$

Cevap: A